

CONTINUATION APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: FATTY ACID DESATURASES AND MUTANT
SEQUENCES THEREOF

APPLICANT: LORIN R. DEBONTE, PH.D., ZHEGONG FAN AND GUO-
HUA MIAO

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL639959554US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

1-29-01
Date of Deposit


Signature

Vince Defante

Typed or Printed Name of Person Signing Certificate

07'48-063003-012901

FATTY ACID DESATURASES AND MUTANT SEQUENCES THEREOF5 Cross-Reference to Related Application

This application is a continuation-in-part of PCT Application Serial No. PCT/US96/20090, filed December 13, 1996, which is a continuation-in-part of U.S. Application Serial No. 08/572,027, filed December 14, 1995.

10 Technical Field

This invention relates to fatty acid desaturases and nucleic acids encoding desaturase proteins. More particularly, the invention relates to nucleic acids encoding delta-12 and delta-15 fatty acid desaturase
15 proteins that affect fatty acid composition in plants, polypeptides produced from such nucleic acids and plants expressing such nucleic acids.

Background of the Invention

Many breeding studies have been conducted to improve
20 the fatty acid profile of Brassica varieties. Pleines and Freidt, Fat Sci. Technol., 90(5), 167-171 (1988) describe plant lines with reduced $C_{18:3}$ levels (2.5-5.8%) combined with high oleic content (73-79%). Rakow and McGregor, J. Amer. Oil Chem. Soc., 50, 400-403 (Oct. 1973) discuss
25 problems associated with selecting mutants for linoleic and linolenic acids. In, Can. J. Plant Sci., 68, 509-511 (Apr. 1988) Stellar summer rape producing seed oil with 3% linolenic acid and 28% linoleic acid is disclosed. Roy and Tarr, Z. Pflanzenzuchtg, 95(3), 201-209 (1985) teaches
30 transfer of genes through an interspecific cross from Brassica juncea into Brassica napus resulting in a reconstituted line combining high linoleic with low linolenic acid content. Roy and Tarr, Plant Breeding, 98,

091904-012904

89-96 (1987) discuss prospects for development of *B. napus* L. having improved linolenic and linolenic acid content. European Patent application 323,753 published July 12, 1989 discloses seeds and oils having greater than 79% oleic acid combined with less than 3.5% linolenic acid. Canvin, Can. J. Botany, 43, 63-69 (1965) discusses the effect of temperature on the fatty acid composition of oils from several seed crops including rapeseed.

Mutations typically are induced with extremely high doses of radiation and/or chemical mutagens (Gaul, H. Radiation Botany (1964) 4:155-232). High dose levels which exceed LD50, and typically reach LD90, led to maximum achievable mutation rates. In mutation breeding of *Brassica* varieties high levels of chemical mutagens alone or combined with radiation have induced a limited number of fatty acid mutations (Rakow, G.Z. Pflanzenzuchtg (1973) 69:62-82). The low α -linolenic acid mutation derived from the Rakow mutation breeding program did not have direct commercial application because of low seed yield. The first commercial cultivar using the low α -linolenic acid mutation derived in 1973 was released in 1988 as the variety Stellar (Scarth, R. et al., Can. J. Plant Sci. (1988) 68:509-511). Stellar was 20% lower yielding than commercial cultivars at the time of its release.

Alterations in fatty acid composition of vegetable oils is desirable for meeting specific food and industrial uses. For example, *Brassica* varieties with increased monounsaturate levels (oleic acid) in the seed oil, and products derived from such oil, would improve lipid nutrition. Canola lines which are low in polyunsaturated fatty acids and high in oleic acid tend to have higher oxidative stability, which is a useful trait for the retail food industry.

Delta-12 fatty acid desaturase (also known as oleic desaturase) is involved in the enzymatic conversion of oleic acid to linoleic acid. Delta-15 fatty acid desaturase (also known as linoleic acid desaturase) is involved in the enzymatic conversion of linoleic acid to α -linolenic acid. A microsomal delta-12 desaturase has been cloned and characterized using T-DNA tagging. Okuley, et al., Plant Cell 6:147-158 (1994). The nucleotide sequences of higher plant genes encoding microsomal delta-12 fatty acid desaturase are described in Lightner et al., WO94/11516. Sequences of higher plant genes encoding microsomal and plastid delta-15 fatty acid desaturases are disclosed in Yadav, N., et al., Plant Physiol., 103:467-476 (1993), WO 93/11245 and Arondel, V. et al., Science, 258:1353-1355 (1992). However, there are no teachings that disclose mutations in delta-12 or delta-15 fatty acid desaturase coding sequences from plants. There is a need in the art for more efficient methods to develop plant lines that contain delta-12 or delta-15 fatty acid desaturase gene sequence mutations effective for altering the fatty acid composition of seeds.

Summary of the Invention

The invention comprises *Brassicaceae* or *Helianthus* seeds, plants and plant lines having at least one mutation that controls the levels of unsaturated fatty acids in plants. One embodiment of the invention is an isolated nucleic acid fragment comprising a nucleotide sequence encoding a mutation from a mutant delta-12 fatty acid desaturase conferring altered fatty composition in seeds when the fragment is present in a plant. A preferred sequence comprises a mutant sequence as shown in Fig. 2. Another embodiment of the invention is an isolated nucleic

acid fragment comprising a nucleotide sequence encoding a mutation from a mutant delta-15 fatty acid desaturase. A plant in this embodiment may be soybean, oilseed *Brassica* species, sunflower, castor bean or corn. The mutant
5 sequence may be derived from, for example, a *Brassica napus*, *Brassica rapa*, *Brassica juncea* or *Helianthus* delta-12 or delta-15 desaturase gene.

Another embodiment of the invention involves a method of producing a *Brassicaceae* or *Helianthus* plant line
10 comprising the steps of: (a) inducing mutagenesis in cells of a starting variety of a *Brassicaceae* or *Helianthus* species; (b) obtaining progeny plants from the mutagenized cells; (c) identifying progeny plants that contain a mutation in a delta-12 or delta-15 fatty acid desaturase
15 gene; and (d) producing a plant line by selfing or crossing. The resulting plant line may be subjected to mutagenesis in order to obtain a line having both a delta-12 desaturase mutation and a delta-15 desaturase mutation.

Yet another embodiment of the invention involves a
20 method of producing plant lines containing altered fatty acid composition comprising: (a) crossing a first plant with a second plant having a mutant delta-12 or delta-15 fatty acid desaturase; (b) obtaining seeds from the cross of step (a); (c) growing fertile plants from such seeds; (d)
25 obtaining progeny seed from the plants of step (c); and (e) identifying those seeds among the progeny that have altered fatty acid composition. Suitable plants are soybean, rapeseed, sunflower, safflower, castor bean and corn. Preferred plants are rapeseed and sunflower.

30 The invention is also embodied in vegetable oil obtained from plants disclosed herein, which vegetable oil has an altered fatty acid composition.

2023年11月11日

5 SEQ ID NO:3 shows a hypothetical DNA sequence of a *Brassica* Fad2 gene having a mutation at nucleotide 316. SEQ ID NO:4 is the deduced amino acid sequence of SEQ ID NO:3.

SEQ ID NO:7 shows a hypothetical DNA sequence of a *Brassica* Fad2 gene having a mutation at nucleotide 515. SEQ ID NO:8 is the deduced amino acid sequence of SEQ ID NO:7.

SEQ ID NO:11 shows the DNA sequence for the coding region of the IMC 129 mutant *Brassica* Fad2-D gene. SEQ ID NO:12 is the deduced amino acid sequence for SEQ ID NO:11.

SEQ ID NO:15 shows the DNA sequence for the coding region of the Q508 mutant *Brassica* Fad2-F gene. SEQ ID NO:16 is the deduced amino acid sequence for SEQ ID NO:15.

Brief Description of the Figures

- 5 -

labeled WSGA 1A represents the $C_{18:1}$ content of the Westar parent. The bar labeled Q508 represents the $C_{18:1}$ content of the Q508 parent.

Sub
C1
5 Figure 2 shows the nucleotide sequences for a
Brassica Fad2-D wild type gene (Fad2-D wt), IMC129 mutant
gene (Fad2-D GA316 IMC129), Fad2-F wild type gene (Fad2-F
wt), Q508 mutant gene (Fad2-F TA515 Q508) and Q4275 mutant
gene (Fad2-F GA908 Q4275).

10 Figure 3 shows the deduced amino acid sequences for
the polynucleotides of Figure 2.

Description of the Preferred Embodiments

All percent fatty acids herein are percent by weight
of the oil of which the fatty acid is a component.

As used herein, a "line" is a group of plants that
15 display little or no genetic variation between individuals
for at least one trait. Such lines may be created by
several generations of self-pollination and selection, or
vegetative propagation from a single parent using tissue or
cell culture techniques. As used herein, the term "variety"
20 refers to a line which is used for commercial production.

The term "mutagenesis" refers to the use of a
mutagenic agent to induce random genetic mutations within a
population of individuals. The treated population, or a
subsequent generation of that population, is then screened
25 for usable trait(s) that result from the mutations. A
"population" is any group of individuals that share a common
gene pool. As used herein " M_0 " is untreated seed. As used
herein, " M_1 " is the seed (and resulting plants) exposed to a
mutagenic agent, while " M_2 " is the progeny (seeds and
30 plants) of self-pollinated M_1 plants, " M_3 " is the progeny of
self-pollinated M_2 plants, and " M_4 " is the progeny of self-
pollinated M_3 plants. " M_5 " is the progeny of self-

pollinated M_4 plants. " M_6 ", " M_7 ", etc. are each the progeny of self-pollinated plants of the previous generation. The term "selfed" as used herein means self-pollinated.

"Stability" or "stable" as used herein means that
5 with respect to a given fatty acid component, the component is maintained from generation to generation for at least two generations and preferably at least three generations at substantially the same level, e.g., preferably $\pm 5\%$. The method of invention is capable of creating lines with
10 improved fatty acid compositions stable up to $\pm 5\%$ from generation to generation. The above stability may be affected by temperature, location, stress and time of planting. Thus, comparison of fatty acid profiles should be made from seeds produced under similar growing conditions.
15 Stability may be measured based on knowledge of prior generation.

Intensive breeding has produced *Brassica* plants whose seed oil contains less than 2% erucic acid. The same varieties have also been bred so that the defatted meal
20 contains less than 30 μmol glucosinolates/gram. "Canola" as used herein refers to plant variety seed or oil which contains less than 2% erucic acid ($C_{22:1}$), and meal with less than 30 μmol glucosinolates/gram.

Applicants have discovered plants with mutations in
25 a delta-12 fatty acid desaturase gene. Such plants have useful alterations in the fatty acid compositions of the seed oil. Such mutations confer, for example, an elevated oleic acid content, a decreased, stabilized linoleic acid content, or both elevated oleic acid and decreased,
30 stabilized linoleic acid content.

Applicants have further discovered plants with mutations in a delta-15 fatty acid desaturase gene. Such plants have useful alterations in the fatty acid composition

of the seed oil, e.g., a decreased, stabilized level of α -linolenic acid.

Applicants have further discovered isolated nucleic acid fragments (polynucleotides) comprising sequences that carry mutations within the coding sequence of delta-12 or delta-15 fatty acid desaturases. The mutations confer desirable alterations in fatty acid levels in the seed oil of plants carrying such mutations. Delta-12 fatty acid desaturase is also known as omega-6 fatty acid desaturase and is sometimes referred to herein as Fad2 or 12-DES. Delta-15 fatty acid desaturase is also known ^{as} ~~on~~ omega-3 fatty acid desaturase and is sometimes referred to herein as Fad3 or 15-DES.

A nucleic acid fragment of the invention may be in the form of RNA or in the form of DNA, including cDNA, synthetic DNA or genomic DNA. The DNA may be double-stranded or single-stranded, and if single-stranded, can be either the coding strand or non-coding strand. An RNA analog may be, for example, mRNA or a combination of ribo- and deoxyribonucleotides. Illustrative examples of a nucleic acid fragment of the invention are the mutant sequences shown in Fig. 3.

A nucleic acid fragment of the invention contains a mutation in a microsomal delta-12 fatty acid desaturase coding sequence or a mutation in a microsomal delta-15 fatty acid desaturase coding sequence. Such a mutation renders the resulting desaturase gene product non-functional in plants, relative to the function of the gene product encoded by the wild-type sequence. The non-functionality of the delta-12 desaturase gene product can be inferred from the decreased level of reaction product (linoleic acid) and increased level of substrate (oleic acid) in plant tissues expressing the mutant sequence, compared to the

corresponding levels in plant tissues expressing the wild-type sequence. The non-functionality of the delta-15 desaturase gene product can be inferred from the decreased level of reaction product (α -linolenic acid) and the increased level of substrate (linoleic acid) in plant tissues expressing the mutant sequence, compared to the corresponding levels in plant tissues expressing the wild-type sequence.

A nucleic acid fragment of the invention may comprise a portion of the coding sequence, e.g., at least about 10 nucleotides, provided that the fragment contains at least one mutation in the coding sequence. The length of a desired fragment depends upon the purpose for which the fragment will be used, e.g., PCR primer, site-directed mutagenesis and the like. In one embodiment, a nucleic acid fragment of the invention comprises the full length coding sequence of a mutant delta-12 or mutant delta-15 fatty acid desaturase, e.g., the mutant sequences of Fig. 3. In other embodiments, a nucleic acid fragment is about 20 to about 50 nucleotides (or base pairs, bp), or about 50 to about 500 nucleotides, or about 500 to about 1200 nucleotides in length.

A mutation in a nucleic acid fragment of the invention may be in any portion of the coding sequence that renders the resulting gene product non-functional. Suitable types of mutations include, without limitation, insertions of nucleotides, deletions of nucleotides, or transitions and transversions in the wild-type coding sequence. Such mutations result in insertions of one or more amino acids, deletions of one or more amino acids, and non-conservative amino acid substitutions in the corresponding gene product. In some embodiments, the sequence of a nucleic acid fragment

may comprise more than one mutation or more than one type of mutation.

Insertion or deletion of amino acids in a coding sequence may, for example, disrupt the conformation of essential alpha-helical or beta-pleated sheet regions of the resulting gene product. Amino acid insertions or deletions may also disrupt binding or catalytic sites important for gene product activity. It is known in the art that the insertion or deletion of a larger number of contiguous amino acids is more likely to render the gene product non-functional, compared to a smaller number of inserted or deleted amino acids.

Non-conservative amino acid substitutions may replace an amino acid of one class with an amino acid of a different class. Non-conservative substitutions may make a substantial change in the charge or hydrophobicity of the gene product. Non-conservative amino acid substitutions may also make a substantial change in the bulk of the residue side chain, e.g., substituting an alanyl residue for a isoleucyl residue.

Examples of non-conservative substitutions include the substitution of a basic amino acid for a non-polar amino acid, or a polar amino acid for an acidic amino acid. Because there are only 20 amino acids encoded in a gene, substitutions that result in a non-functional gene product may be determined by routine experimentation, incorporating amino acids of a different class in the region of the gene product targeted for mutation.

Preferred mutations are in a region of the nucleic acid encoding an amino acid sequence motif that is conserved among delta-12 fatty acid desaturases or delta-15 fatty acid desaturases, such as a His-Xaa-Xaa-Xaa-His motif (Tables 1-3). An example of a suitable region has a conserved HECGH

Sub
C2

Sub
C2
cancel

A

5 motif that is found, for example, in nucleotides
corresponding to amino acids 105 to 109 of the *Arabidopsis*
and *Brassica* delta-12 desaturase sequences, in nucleotides
corresponding to amino acids 101 to 105 of the soybean
delta-12 desaturase sequence and in nucleotides
corresponding to amino acids 111 to 115 of the maize delta-
12 desaturase sequence. See e.g., WO 94/11516¹¹⁵¹⁶; Okuley et
al., *Plant Cell* 6:147-158 (1994). The one letter amino acid
designations used herein are described in Alberts, B. et
10 al., *Molecular Biology of the Cell*, 3rd edition, Garland
Publishing, New York, 1994. Amino acids flanking this motif
are also highly conserved among delta-12 and delta-15
desaturases and are also suitable candidates for mutations
in fragments of the invention.

15 An illustrative embodiment of a mutation in a
nucleic acid fragment of the invention is a Glu to Lys
substitution in the HECGH motif of a *Brassica* microsomal
delta-12 desaturase sequence, either the D form or the F
form. This mutation results in the sequence HECGH being
20 changed to HECGH as seen by comparing SEQ ID NO:10 (wild-
type D form) to SEQ ID NO:12 (mutant D form). A similar
mutation in other Fad-2 sequences is contemplated to result
in a non-functional gene product. (Compare SEQ ID NO:2 to
SEQ ID NO:4)

25 A similar motif may be found at amino acids 101 to
105 of the *Arabidopsis* microsomal delta-15 fatty acid
desaturase, as well as in the corresponding rape and soybean
desaturases (Table 5). See, e.g., WO 93/11245; Arondel, V.
et al., *Science*, 258:1153-1155 (1992); Yadav, N. et al.,
30 *Plant Physiol.*, 103:467-476 (1993). Plastid delta-15 fatty
acids have a similar motif (Table 5).

Sub
C3

Among the types of mutations in an HECGH motif that
render the resulting gene product non-functional are non-

Sub
C3
C4
C5
C6
C7
C8
C9
C10
C11
C12
C13
C14
C15
C16
C17
C18
C19
C20
C21
C22
C23
C24
C25
C26
C27
C28
C29
C30
C31
C32
C33
C34
C35
C36
C37
C38
C39
C40
C41
C42
C43
C44
C45
C46
C47
C48
C49
C50
C51
C52
C53
C54
C55
C56
C57
C58
C59
C60
C61
C62
C63
C64
C65
C66
C67
C68
C69
C70
C71
C72
C73
C74
C75
C76
C77
C78
C79
C80
C81
C82
C83
C84
C85
C86
C87
C88
C89
C90
C91
C92
C93
C94
C95
C96
C97
C98
C99
C100
C101
C102
C103
C104
C105
C106
C107
C108
C109
C110
C111
C112
C113
C114
C115
C116
C117
C118
C119
C120
C121
C122
C123
C124
C125
C126
C127
C128
C129
C130
C131
C132
C133
C134
C135
C136
C137
C138
C139
C140
C141
C142
C143
C144
C145
C146
C147
C148
C149
C150
C151
C152
C153
C154
C155
C156
C157
C158
C159
C160
C161
C162
C163
C164
C165
C166
C167
C168
C169
C170
C171
C172
C173
C174
C175
C176
C177
C178
C179
C180
C181
C182
C183
C184
C185
C186
C187
C188
C189
C190
C191
C192
C193
C194
C195
C196
C197
C198
C199
C200
C201
C202
C203
C204
C205
C206
C207
C208
C209
C210
C211
C212
C213
C214
C215
C216
C217
C218
C219
C220
C221
C222
C223
C224
C225
C226
C227
C228
C229
C230
C231
C232
C233
C234
C235
C236
C237
C238
C239
C240
C241
C242
C243
C244
C245
C246
C247
C248
C249
C250
C251
C252
C253
C254
C255
C256
C257
C258
C259
C260
C261
C262
C263
C264
C265
C266
C267
C268
C269
C270
C271
C272
C273
C274
C275
C276
C277
C278
C279
C280
C281
C282
C283
C284
C285
C286
C287
C288
C289
C290
C291
C292
C293
C294
C295
C296
C297
C298
C299
C300
C301
C302
C303
C304
C305
C306
C307
C308
C309
C310
C311
C312
C313
C314
C315
C316
C317
C318
C319
C320
C321
C322
C323
C324
C325
C326
C327
C328
C329
C330
C331
C332
C333
C334
C335
C336
C337
C338
C339
C340
C341
C342
C343
C344
C345
C346
C347
C348
C349
C350
C351
C352
C353
C354
C355
C356
C357
C358
C359
C360
C361
C362
C363
C364
C365
C366
C367
C368
C369
C370
C371
C372
C373
C374
C375
C376
C377
C378
C379
C380
C381
C382
C383
C384
C385
C386
C387
C388
C389
C390
C391
C392
C393
C394
C395
C396
C397
C398
C399
C400
C401
C402
C403
C404
C405
C406
C407
C408
C409
C410
C411
C412
C413
C414
C415
C416
C417
C418
C419
C420
C421
C422
C423
C424
C425
C426
C427
C428
C429
C430
C431
C432
C433
C434
C435
C436
C437
C438
C439
C440
C441
C442
C443
C444
C445
C446
C447
C448
C449
C450
C451
C452
C453
C454
C455
C456
C457
C458
C459
C460
C461
C462
C463
C464
C465
C466
C467
C468
C469
C470
C471
C472
C473
C474
C475
C476
C477
C478
C479
C480
C481
C482
C483
C484
C485
C486
C487
C488
C489
C490
C491
C492
C493
C494
C495
C496
C497
C498
C499
C500
C501
C502
C503
C504
C505
C506
C507
C508
C509
C510
C511
C512
C513
C514
C515
C516
C517
C518
C519
C520
C521
C522
C523
C524
C525
C526
C527
C528
C529
C530
C531
C532
C533
C534
C535
C536
C537
C538
C539
C540
C541
C542
C543
C544
C545
C546
C547
C548
C549
C550
C551
C552
C553
C554
C555
C556
C557
C558
C559
C560
C561
C562
C563
C564
C565
C566
C567
C568
C569
C570
C571
C572
C573
C574
C575
C576
C577
C578
C579
C580
C581
C582
C583
C584
C585
C586
C587
C588
C589
C590
C591
C592
C593
C594
C595
C596
C597
C598
C599
C600
C601
C602
C603
C604
C605
C606
C607
C608
C609
C610
C611
C612
C613
C614
C615
C616
C617
C618
C619
C620
C621
C622
C623
C624
C625
C626
C627
C628
C629
C630
C631
C632
C633
C634
C635
C636
C637
C638
C639
C640
C641
C642
C643
C644
C645
C646
C647
C648
C649
C650
C651
C652
C653
C654
C655
C656
C657
C658
C659
C660
C661
C662
C663
C664
C665
C666
C667
C668
C669
C670
C671
C672
C673
C674
C675
C676
C677
C678
C679
C680
C681
C682
C683
C684
C685
C686
C687
C688
C689
C690
C691
C692
C693
C694
C695
C696
C697
C698
C699
C700
C701
C702
C703
C704
C705
C706
C707
C708
C709
C710
C711
C712
C713
C714
C715
C716
C717
C718
C719
C720
C721
C722
C723
C724
C725
C726
C727
C728
C729
C730
C731
C732
C733
C734
C735
C736
C737
C738
C739
C740
C741
C742
C743
C744
C745
C746
C747
C748
C749
C750
C751
C752
C753
C754
C755
C756
C757
C758
C759
C760
C761
C762
C763
C764
C765
C766
C767
C768
C769
C770
C771
C772
C773
C774
C775
C776
C777
C778
C779
C780
C781
C782
C783
C784
C785
C786
C787
C788
C789
C790
C791
C792
C793
C794
C795
C796
C797
C798
C799
C800
C801
C802
C803
C804
C805
C806
C807
C808
C809
C810
C811
C812
C813
C814
C815
C816
C817
C818
C819
C820
C821
C822
C823
C824
C825
C826
C827
C828
C829
C830
C831
C832
C833
C834
C835
C836
C837
C838
C839
C840
C841
C842
C843
C844
C845
C846
C847
C848
C849
C850
C851
C852
C853
C854
C855
C856
C857
C858
C859
C860
C861
C862
C863
C864
C865
C866
C867
C868
C869
C870
C871
C872
C873
C874
C875
C876
C877
C878
C879
C880
C881
C882
C883
C884
C885
C886
C887
C888
C889
C890
C891
C892
C893
C894
C895
C896
C897
C898
C899
C900
C901
C902
C903
C904
C905
C906
C907
C908
C909
C910
C911
C912
C913
C914
C915
C916
C917
C918
C919
C920
C921
C922
C923
C924
C925
C926
C927
C928
C929
C930
C931
C932
C933
C934
C935
C936
C937
C938
C939
C940
C941
C942
C943
C944
C945
C946
C947
C948
C949
C950
C951
C952
C953
C954
C955
C956
C957
C958
C959
C960
C961
C962
C963
C964
C965
C966
C967
C968
C969
C970
C971
C972
C973
C974
C975
C976
C977
C978
C979
C980
C981
C982
C983
C984
C985
C986
C987
C988
C989
C990
C991
C992
C993
C994
C995
C996
C997
C998
C999
C1000
C1001
C1002
C1003
C1004
C1005
C1006
C1007
C1008
C1009
C1010
C1011
C1012
C1013
C1014
C1015
C1016
C1017
C1018
C1019
C1020
C1021
C1022
C1023
C1024
C1025
C1026
C1027
C1028
C1029
C1030
C1031
C1032
C1033
C1034
C1035
C1036
C1037
C1038
C1039
C1040
C1041
C1042
C1043
C1044
C1045
C1046
C1047
C1048
C1049
C1050
C1051
C1052
C1053
C1054
C1055
C1056
C1057
C1058
C1059
C1060
C1061
C1062
C1063
C1064
C1065
C1066
C1067
C1068
C1069
C1070
C1071
C1072
C1073
C1074
C1075
C1076
C1077
C1078
C1079
C1080
C1081
C1082
C1083
C1084
C1085
C1086
C1087
C1088
C1089
C1090
C1091
C1092
C1093
C1094
C1095
C1096
C1097
C1098
C1099
C1100
C1101
C1102
C1103
C1104
C1105
C1106
C1107
C1108
C1109
C1110
C1111
C1112
C1113
C1114
C1115
C1116
C1117
C1118
C1119
C1120
C1121
C1122
C1123
C1124
C1125
C1126
C1127
C1128
C1129
C1130
C1131
C1132
C1133
C1134
C1135
C1136
C1137
C1138
C1139
C1140
C1141
C1142
C1143
C1144
C1145
C1146
C1147
C1148
C1149
C1150
C1151
C1152
C1153
C1154
C1155
C1156
C1157
C1158
C1159
C1160
C1161
C1162
C1163
C1164
C1165
C1166
C1167
C1168
C1169
C1170
C1171
C1172
C1173
C1174
C1175
C1176
C1177
C1178
C1179
C1180
C1181
C1182
C1183
C1184
C1185
C1186
C1187
C1188
C1189
C1190
C1191
C1192
C1193
C1194
C1195
C1196
C1197
C1198
C1199
C1200
C1201
C1202
C1203
C1204
C1205
C1206
C1207
C1208
C1209
C1210
C1211
C1212
C1213
C1214
C1215
C1216
C1217
C1218
C1219
C1220
C1221
C1222
C1223
C1224
C1225
C1226
C1227
C1228
C1229
C1230
C1231
C1232
C1233
C1234
C1235
C1236
C1237
C1238
C1239
C1240
C1241
C1242
C1243
C1244
C1245
C1246
C1247
C1248
C1249
C1250
C1251
C1252
C1253
C1254
C1255
C1256
C1257
C1258
C1259
C1260
C1261
C1262
C1263
C1264
C1265
C1266
C1267
C1268
C1269
C1270
C1271
C1272
C1273
C1274
C1275
C1276
C1277
C1278
C1279
C1280
C1281
C1282
C1283
C1284
C1285
C1286
C1287
C1288
C1289
C1290
C1291
C1292
C1293
C1294
C1295
C1296
C1297
C1298
C1299
C1300
C1301
C1302
C1303
C1304
C1305
C1306
C1307
C1308
C1309
C1310
C1311
C1312
C1313
C1314
C1315
C1316
C1317
C1318
C1319
C1320
C1321
C1322
C1323
C1324
C1325
C1326
C1327
C1328
C1329
C1330
C1331
C1332
C1333
C1334
C1335
C1336
C1337
C1338
C1339
C1340
C1341
C1342
C1343
C1344
C1345
C1346
C1347
C1348
C1349
C1350
C1351
C1352
C1353
C1354
C1355
C1356
C1357
C1358
C1359
C1360
C1361
C1362
C1363
C1364
C1365
C1366
C1367
C1368
C1369
C1370
C1371
C1372
C1373
C1374
C1375
C1376
C1377
C1378
C1379
C1380
C1381
C1382
C1383
C1384
C1385
C1386
C1387
C1388
C1389
C1390
C1391
C1392
C1393
C1394
C1395
C1396
C1397
C1398
C1399
C1400
C1401
C1402
C1403
C1404
C1405
C1406
C1407
C1408
C1409
C1410
C1411
C1412
C1413
C1414
C1415
C1416
C1417
C1418
C1419
C1420
C1421
C1422
C1423
C1424
C1425
C1426
C1427
C1428
C1429
C1430
C1431
C1432
C1433
C1434
C1435
C1436
C1437
C1438
C1439
C1440
C1441
C1442
C1443
C1444
C1445
C1446
C1447
C1448
C1449
C1450
C1451
C1452
C1453
C1454
C1455
C1456
C1457
C1458
C1459
C1460
C1461
C1462
C1463
C1464
C1465
C1466
C1467
C1468
C1469
C1470
C1471
C1472
C1473
C1474
C1475
C1476
C1477
C1478
C1479
C1480
C1481
C1482
C1483
C1484
C1485
C1486
C1487
C1488
C1489
C1490
C1491
C1492
C1493
C1494
C1495
C1496
C1497
C1498
C1499
C1500
C1501
C1502
C1503
C1504
C1505
C1506
C1507
C1508
C1509
C1510
C1511
C1512
C1513
C1514
C1515
C1516
C1517
C1518
C1519
C1520
C1521
C1522
C1523
C1524
C1525
C1526
C1527
C1528
C1529
C1530
C1531
C1532
C1533
C1534
C1535
C1536
C1537
C1538
C1539
C1540
C1541
C1542
C1543
C1544
C1545
C1546
C1547
C1548
C1549
C1550
C1551
C1552
C1553
C1554
C1555
C1556
C1557
C1558
C1559
C1560
C1561
C1562
C1563
C1564
C1565
C1566
C1567
C1568
C1569
C1570
C1571
C1572
C1573
C1574
C1575
C1576
C1577
C1578
C1579
C1580
C1581
C1582
C1583
C1584
C1585
C1586
C1587
C1588
C1589
C1590
C1591
C1592
C1593
C1594
C1595
C1596
C1597
C1598
C1599
C1600
C1601
C1602
C1603
C1604
C1605
C1606
C1607
C1608
C1609
C1610
C1611
C1612
C1613
C1614
C1615
C1616
C1617
C1618
C1619
C1620
C1621
C1622
C1623
C1624
C1625
C1626
C1627
C1628
C1629
C1630
C1631
C1632
C1633
C1634
C1635
C1636
C1637
C1638
C1639
C1640
C1641
C1642
C1643
C1644
C1645
C1646
C1647
C1648
C1649
C1650
C1651
C1652
C1653
C1654
C1655
C1656
C1657
C1658
C1659
C1660
C1661
C1662
C1663
C1664
C1665
C1666
C1667
C1668
C1669
C1670
C1671
C1672
C1673
C1674
C1675
C1676
C1677
C1678
C1679
C1680
C1681
C1682
C1683
C1684
C1685
C1686
C1687
C1688
C1689
C1690
C1691
C1692
C1693
C1694
C1695
C1696
C1697
C1698
C1699
C1700
C1701
C1702
C1703
C1704
C1705
C1706
C1707
C1708
C1709
C1710
C1711
C1712
C1713
C1714
C1715
C1716
C1717
C1718
C1719
C1720
C1721
C1722
C1723
C1724
C1725
C1726
C1727
C1728
C1729
C1730
C1731
C1732
C1733
C1734
C1735
C1736
C1737
C1738
C1739
C1740
C1741
C1742
C1743
C1744
C1745
C1746
C1747
C1748
C1749
C1750
C1751
C1752
C1753
C1754
C1755
C1756
C1757
C1758
C1759
C1760
C1761
C1762
C1763
C1764
C1765
C1766
C1767
C1768
C1769
C1770
C1771
C1772
C1773
C1774
C1775
C1776
C1777
C1778
C1779
C1780
C1781
C1782
C1783
C1784
C1785
C1786
C1787
C1788
C1789
C1790
C1791
C1792
C1793
C1794
C1795
C1796
C1797
C1798
C1799
C1800
C1801
C1802
C1803
C1804
C1805
C1806
C1807
C1808
C1809
C1810
C1811
C1812
C1813
C1814
C1815
C1816
C1817
C1818
C1819
C1820
C1821
C1822
C1823
C1824
C1825
C1826
C1827
C1828
C1829
C1830
C1831
C1832
C1833
C1834
C1835
C1836
C1837
C1838
C1839
C1840
C1841
C1842
C1843
C1844
C1845
C1846
C1847
C1848
C1849
C1850
C1851
C1852
C1853
C1854
C1855
C1856
C1857
C1858
C1859
C1860
C1861
C1862
C1863
C1864
C1865
C1866
C1867
C1868
C1869
C1870
C1871
C1872
C1873
C1874
C1875
C1876
C1877
C1878
C1879
C1880
C1881
C1882
C1883
C1884
C1885
C1886
C1887
C1888
C1889
C1890
C1891
C1892
C1893
C1894
C1895
C1896
C1897
C1898
C1899
C1900
C1901
C1902
C1903
C1904
C1905
C1906
C1907
C1908
C1909
C1910
C1911
C1912
C1913
C1914
C1915
C1916
C1917
C1918
C1919
C1920
C1921
C1922
C1923
C1924

TABLE 1

Alignment of Amino Acid Sequences from Microsomal
Delta-12 Fatty Acid Desaturases

Sub
CS >

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>Arabidopsis thaliana</i>	100-129	IWVIAHECGH HAFSDYQWLD DTVGLIFHSF
	<i>Glycine max</i>	96-125	VWVIAHECGH HAFSKYQWVD DVVGLTLHST
	<i>Zea mays</i>	106-135	VWVIAHECGH HAFSDYSLLD DVVGLVLHSS
	<i>Ricinus communis</i> ^a	1- 29	WVMAHDCGH HAFSDYQLLD DVVGLILHSC
	<i>Brassica napus D</i>	100-128	VWVIAHECGH HAFSDYQWLD DTVGLIFHS
10	<i>Brassica napus F</i>	100-128	VWVIAHECGH HAFSDYQWLD DTVGLIFHS

^a from plasmid pRF2-1C

TABLE 2

Alignment of Amino Acid Sequences from Microsomal
Delta-12 Fatty Acid Desaturases

09771904-012601

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
15	<i>Arabidopsis thaliana</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Glycine max</i>	126-154	LLVPYFSWKI SHRRHHSNTG SLDRDEVFV
	<i>Zea mays</i>	136-164	LMVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Ricinus communis</i> ^a	30- 58	LLVPYFSWKH SHRRHHSNTG SLERDEVFV
20	<i>Brassica napus D</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Brassica napus F</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV

^a from plasmid pRF2-1C

TABLE 3

Alignment of Amino Acid Sequences from Microsomal
Delta-12 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
25	<i>Arabidopsis thaliana</i>	298-333	DRDYGILNKV FHNITDTHVA HHLFSTMPHY NAMEAT
	<i>Glycine max</i>	294-329	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT
	<i>Zea mays</i>	305-340	DRDYGILNRV FHNITDTHVA HHLFSTMPHY HAMEAT
30	<i>Ricinus communis</i> ^a	198-224	DRDYGILNKV FHNITDTQVA HHLF TMP
	<i>Brassica napus D</i>	299-334	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT
	<i>Brassica napus F</i>	299-334	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT

^a from plasmid pRF2-1C

TABLE 4

Alignment of Conserved Amino Acids from Microsomal
Delta-12 Fatty Acid Desaturases

Sub
C5
amald

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>Arabidopsis thaliana</i>	165-180	IKWYGKYLNN PLGRIM
	<i>Glycine max</i>	161-176	VAWFSLYLNN PLGRAV
	<i>Zea mays</i>	172-187	PWYTPYVYNN PVGRVV
	<i>Ricinus communis</i> ^a	65-80	IRWYSKYLNN PPGRIM
	<i>Brassica napus</i> D	165-180	IKWYGKYLNN PLGRTV
10	<i>Brassica napus</i> F	165-180	IKWYGKYLNN PLGRTV

^a from plasmid pRF2-1C

TABLE 5

Alignment of Conserved Amino Acids from Plastid and Microsomal
Delta-15 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
15	<i>Arabidopsis thaliana</i> ^a	156-177	WALFVLGHD CGHGSFSNDP KLN
	<i>Brassica napus</i> ^a	114-135	WALFVLGHD CGHGSFSNDP RLN
	<i>Glycine max</i> ^a	164-185	WALFVLGHD CGHGSFSNNS KLN
	<i>Arabidopsis thaliana</i>	94-115	WAIFVLGHD CGHGSFSDIP LLN
20	<i>Brassica napus</i>	87-109	WALFVLGHD CGHGSFSNDP RLN
	<i>Glycine max</i>	93-114	WALFVLGHD CGHGSFSDSP PLN

^a Plastid sequences

TABLE 6

Alignment of Conserved Amino Acids from Plastid and Microsomal
Delta-15 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
25	<i>A. thaliana</i> ^a	188-216	ILVPYHGWRI SHRTHHQNHG HVENDESWH
	<i>B. napus</i> ^a	146-174	ILVPYHGWRI SHRTHHQNHG HVENDESWH
	<i>Glycine max</i> ^a	196-224	ILVPYHGWRI SHRTHHQNHG HAENDESWH
30	<i>A. thaliana</i>	126-154	ILVPYHGWRI SHRTHHQNHG HVENDESWV
	<i>Brassica napus</i>	117-145	ILVPYHGWRI SHRTHHQNHG HVENDESWV
	<i>Glycine max</i>	125-153	ILVPYHGWRI SHRTHHQNHG HIEKDESWV

^a Plastid sequences

The conservation of amino acid motifs and their
relative positions indicates that regions of a delta-12 or

delta-15 fatty acid desaturase that can be mutated in one species to generate a non-functional desaturase can be mutated in the corresponding region from other species to generate a non-functional delta-12 desaturase or delta-15 desaturase gene product in that species.

Mutations in any of the regions of Tables 1-6 are specifically included within the scope of the invention and are substantially identical to those mutations exemplified herein, provided that such mutation (or mutations) renders the resulting desaturase gene product non-functional, as discussed hereinabove.

A nucleic acid fragment containing a mutant sequence can be generated by techniques known to the skilled artisan. Such techniques include, without limitation, site-directed mutagenesis of wild-type sequences and direct synthesis using automated DNA synthesizers.

A nucleic acid fragment containing a mutant sequence can also be generated by mutagenesis of plant seeds or regenerable plant tissue by, e.g., ethyl methane sulfonate, X-rays or other mutagens. With mutagenesis, mutant plants having the desired fatty acid phenotype in seeds are identified by known techniques and a nucleic acid fragment containing the desired mutation is isolated from genomic DNA or RNA of the mutant line. The site of the specific mutation is then determined by sequencing the coding region of the delta-12 desaturase or delta-15 desaturase gene. Alternatively, labeled nucleic acid probes that are specific for desired mutational events can be used to rapidly screen a mutagenized population.

The disclosed method may be applied to all oilseed Brassica species, and to both Spring and Winter maturing types within each species. Physical mutagens, including but not limited to X-rays, UV rays, and other physical

0974904-012901
treatments which cause chromosome damage, and other chemical mutagens, including but not limited to ethidium bromide, nitrosoguanidine, diepoxybutane etc. may also be used to induce mutations. The mutagenesis treatment may also be applied to other stages of plant development, including but not limited to cell cultures, embryos, microspores and shoot apices.

"Stable mutations" as used herein are defined as M_s or more advanced lines which maintain a selected altered fatty acid profile for a minimum of three generations, including a minimum of two generations under field conditions, and exceeding established statistical thresholds for a minimum of two generations, as determined by gas chromatographic analysis of a minimum of 10 randomly selected seeds bulked together. Alternatively, stability may be measured in the same way by comparing to subsequent generations. In subsequent generations, stability is defined as having similar fatty acid profiles in the seed as that of the prior or subsequent generation when grown under substantially similar conditions.

Mutation breeding has traditionally produced plants carrying, in addition to the trait of interest, multiple, deleterious traits, e.g., reduced plant vigor and reduced fertility. Such traits may indirectly affect fatty acid composition, producing an unstable mutation; and/or reduce yield, thereby reducing the commercial utility of the invention. To eliminate the occurrence of deleterious mutations and reduce the load of mutations carried by the plant, a low mutagen dose is used in the seed treatments to create an LD30 population. This allows for the rapid selection of single gene mutations for fatty acid traits in agronomic backgrounds which produce acceptable yields.

Sub
C6 >
The seeds of several different fatty acid lines have been deposited with the American Type Culture Collection and have the following accession numbers.

	<u>Line</u>	<u>Accession No.</u>	<u>Deposit Date</u>
5	A129.5	40811	May 25, 1990
	A133.1	40812	May 25, 1990
	M3032.1	75021	June 7, 1991
	M3062.8	75025	June 7, 1991
	M3028.10	75026	June 7, 1991
10	IMC130	75446	April 16, 1993
	Q4275	97569	May 10, 1996

09771904-012901
In some plant species or varieties more than one form of endogenous microsomal delta-12 desaturase may be found. In amphidiploids, each form may be derived from one of the parent genomes making up the species under consideration. Plants with mutations in both forms have a fatty acid profile that differs from plants with a mutation in only one form. An example of such a plant is *Brassica napus* line Q508, a doubly-mutagenized line containing a mutant D-form of delta-12 desaturase (SEQ ID NO:11) and a mutant F-form of delta-12 desaturase (SEQ ID NO:15). Another example is line Q4275, which contains a mutant D-form of delta-12 desaturase (SEQ ID NO:11) and a mutant F-form of delta-12 desaturase (SEQ ID NO:17). See Figs. 2-3.

Preferred host or recipient organisms for introduction of a nucleic acid fragment of the invention are the oil-producing species, such as soybean (*Glycine max*), rapeseed (e.g., *Brassica napus*, *B. rapa* and *B. juncea*), sunflower (*Helianthus annuus*), castor bean (*Ricinus communis*), corn (*Zea mays*), and safflower (*Carthamus tinctorius*).

A nucleic acid fragment of the invention may further comprise additional nucleic acids. For example, a nucleic acid encoding a secretory or leader amino acid sequence can

be linked to a mutant desaturase nucleic acid fragment such that the secretory or leader sequence is fused in-frame to the amino terminal end of a mutant delta-12 or delta-15 desaturase polypeptide. Other nucleic acid fragments are known in the art that encode amino acid sequences useful for fusing in-frame to the mutant desaturase polypeptides disclosed herein. See, e.g., U.S. 5,629,193 incorporated herein by reference. A nucleic acid fragment may also have one or more regulatory elements operably linked thereto.

The present invention also comprises nucleic acid fragments that selectively hybridize to mutant desaturase sequences. Such a nucleic acid fragment typically is at least 15 nucleotides in length. Hybridization typically involves Southern analysis (Southern blotting), a method by which the presence of DNA sequences in a target nucleic acid mixture are identified by hybridization to a labeled oligonucleotide or DNA fragment probe. Southern analysis typically involves electrophoretic separation of DNA digests on agarose gels, denaturation of the DNA after electrophoretic separation, and transfer of the DNA to nitrocellulose, nylon, or another suitable membrane support for analysis with a radiolabeled, biotinylated, or enzyme-labeled probe as described in sections 9.37-9.52 of Sambrook et al., (1989) *Molecular Cloning*, second edition, Cold Spring Harbor Laboratory, Plainview; NY.

A nucleic acid fragment can hybridize under moderate stringency conditions or, preferably, under high stringency conditions to a mutant desaturase sequence. High stringency conditions are used to identify nucleic acids that have a high degree of homology to the probe. High stringency conditions can include the use of low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate (0.1X SSC); 0.1% sodium lauryl

09771904-012901

sulfate (SDS) at 50-65°C. Alternatively, a denaturing agent such as formamide can be employed during hybridization, e.g., 50% formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42°C. Another example is the use of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS.

Moderate stringency conditions refers to hybridization conditions used to identify nucleic acids that have a lower degree of identity to the probe than do nucleic acids identified under high stringency conditions. Moderate stringency conditions can include the use of higher ionic strength and/or lower temperatures for washing of the hybridization membrane, compared to the ionic strength and temperatures used for high stringency hybridization. For example, a wash solution comprising 0.060 M NaCl/0.0060 M sodium citrate (4X SSC) and 0.1% sodium lauryl sulfate (SDS) can be used at 50°C, with a last wash in 1X SSC, at 65°C. Alternatively, a hybridization wash in 1X SSC at 37°C can be used.

Hybridization can also be done by Northern analysis (Northern blotting), a method used to identify RNAs that hybridize to a known probe such as an oligonucleotide, DNA fragment, cDNA or fragment thereof, or RNA fragment. The probe is labeled with a radioisotope such as ³²P, by biotinylation or with an enzyme. The RNA to be analyzed can be usually electrophoretically separated on an agarose or polyacrylamide gel, transferred to nitrocellulose, nylon, or other suitable membrane, and hybridized with the probe,

using standard techniques well known in the art such as those described in sections 7.39-7.52 of Sambrook et al., *supra*.

5 A polypeptide of the invention comprises an isolated polypeptide having a mutant amino acid sequence, as well as derivatives and analogs thereof. See, e.g., the mutant amino acid sequences of Fig. 3. By "isolated" is meant a polypeptide that is expressed and produced in an environment other than the environment in which the polypeptide is
10 naturally expressed and produced. For example, a plant polypeptide is isolated when expressed and produced in bacteria or fungi. A polypeptide of the invention also comprises variants of the mutant desaturase polypeptides disclosed herein, as discussed above.

15 In one embodiment of the claimed invention, a plant contains both a delta-12 desaturase mutation and a delta-15 desaturase mutation. Such plants can have a fatty acid composition comprising very high oleic acid and very low alpha-linolenic acid levels. Mutations in delta-12
20 desaturase and delta-15 desaturase may be combined in a plant by making a genetic cross between delta-12 desaturase and delta-15 desaturase single mutant lines. A plant having a mutation in delta-12 fatty acid desaturase is crossed or mated with a second plant having a mutation in delta-15
25 fatty acid desaturase. Seeds produced from the cross are planted and the resulting plants are selfed in order to obtain progeny seeds. These progeny seeds are then screened in order to identify those seeds carrying both mutant genes.

Alternatively, a line possessing either a delta-12
30 desaturase or a delta-15 desaturase mutation can be subjected to mutagenesis to generate a plant or plant line having mutations in both delta-12 desaturase and delta-15 desaturase. For example, the IMC 129 line has a mutation in

the coding region (Glu₁₀₆ to Lys₁₀₆) of the D form of the
microsomal delta-12 desaturase structural gene. Cells
(e.g., seeds) of this line can be mutagenized to induce a
mutation in a delta-15 desaturase gene, resulting in a plant
5 or plant line carrying a mutation in a delta-12 fatty acid
desaturase gene and a mutation in a delta-15 fatty acid
desaturase gene.

Progeny includes descendants of a particular plant
or plant line, e.g., seeds developed on an instant plant are
10 descendants. Progeny of an instant plant include seeds
formed on F₁, F₂, F₃, and subsequent generation plants, or
seeds formed on BC₁, BC₂, BC₃, and subsequent generation
plants.

Plants according to the invention preferably contain
15 an altered fatty acid composition. For example, oil
obtained from seeds of such plants may have from about 69 to
about 90% oleic acid, based on the total fatty acid
composition of the seed. Such oil preferably has from about
74 to about 90% oleic acid, more preferably from about 80 to
20 about 90% oleic acid. In some embodiments, oil obtained
from seeds produced by plants of the invention may have from
about 2.0% to about 5.0% saturated fatty acids, based on
total fatty acid composition of the seeds. In some
embodiments, oil obtained from seeds of the invention may
25 have from about 1.0% to about 14.0% linoleic acid, or from
about 0.5% to about 10.0% α -linolenic acid.

Oil composition typically is analyzed by crushing
and extracting fatty acids from bulk seed samples (e.g., 10
seeds). Fatty acid triglycerides in the seed are hydrolyzed
30 and converted to fatty acid methyl esters. Those seeds
having an altered fatty acid composition may be identified
by techniques known to the skilled artisan, e.g., gas-liquid
chromatography (GLC) analysis of a bulked seed sample or of

5 a single half-seed. Half-seed analysis is well known in the art to be useful because the viability of the embryo is maintained and thus those seeds having a desired fatty acid profile may be planted to from the next generation.

10 However, half-seed analysis is also known to be an inaccurate representation of genotype of the seed being analyzed. Bulk seed analysis typically yields a more accurate representation of the fatty acid profile of a given genotype. Fatty acid composition can also be determined on larger samples, e.g., oil obtained by pilot plant or commercial scale refining, bleaching and deodorizing of endogenous oil in the seeds.

15 The nucleic acid fragments of the invention can be used as markers in plant genetic mapping and plant breeding programs. Such markers may include restriction fragment length polymorphism (RFLP), random amplification polymorphism detection (RAPD), polymerase chain reaction (PCR) or self-sustained sequence replication (3SR) markers, for example. Marker-assisted breeding techniques may be used to identify and follow a desired fatty acid composition during the breeding process. Marker-assisted breeding techniques may be used in addition to, or as an alternative to, other sorts of identification techniques. An example of marker-assisted breeding is the use of PCR primers that specifically amplify a sequence containing a desired mutation in delta-12 desaturase or delta-15 desaturase.

25 Methods according to the invention are useful in that the resulting plants and plant lines have desirable seed fatty acid compositions as well as superior agronomic properties compared to known lines having altered seed fatty acid composition. Superior agronomic characteristics include, for example, increased seed germination percentage, increased seedling vigor, increased resistance to seedling

09271904-012901
T06270-4067260

fungus diseases (damping off, root rot and the like),
increased yield, and improved standability.

While the invention is susceptible to various
modifications and alternative forms, certain specific
5 embodiments thereof are described in the general methods and
examples set forth below. For example the invention may be
applied to all *Brassica* species, including *B. rapa*, *B.*
juncea, and *B. hirta*, to produce substantially similar
results. It should be understood, however, that these
10 examples are not intended to limit the invention to the
particular forms disclosed but, instead the invention is to
cover all modifications, equivalents and alternatives
falling within the scope of the invention. This includes
the use of somaclonal variation; physical or chemical
15 mutagenesis of plant parts; anther, microspore or ovary
culture followed by chromosome doubling; or self- or cross-
pollination to transmit the fatty acid trait, alone or in
combination with other traits, to develop new *Brassica*
lines.

20 EXAMPLE 1

Mutagenesis

Seeds of Westar, a Canadian (*Brassica napus*) spring
canola variety, were subjected to chemical mutagenesis.
Westar is a registered Canadian spring variety with canola
25 quality. The fatty acid composition of field-grown Westar,
3.9% C_{16:0}, 1.9% C_{18:0}, 67.5% C_{18:1}, 17.6% C_{18:2}, 7.4% C_{18:3}, <2%
C_{20:1} + C_{22:1}, has remained stable under commercial
production, with $\pm 10\%$ deviation, since 1982.

Prior to mutagenesis, 30,000 seeds of *B. napus* cv.
30 Westar seeds were preimbibed in 300-seed lots for two hours
on wet filter paper to soften the seed coat. The preimbibed
seeds were placed in 80 mM ethylmethanesulfonate (EMS) for

four hours. Following mutagenesis, the seeds were rinsed three times in distilled water. The seeds were sown in 48-well flats containing Pro-Mix. Sixty-eight percent of the mutagenized seed germinated. The plants were maintained at 25°C/15°C, 14/10 hr day/night conditions in the greenhouse. At flowering, each plant was individually self-pollinated.

M₂ seed from individual plants were individually catalogued and stored, approximately 15,000 M₂ lines was planted in a summer nursery in Carman, Manitoba. The seed from each selfed plant were planted in 3-meter rows with 6-inch row spacing. Westar was planted as the check variety. Selected lines in the field were selfed by bagging the main raceme of each plant. At maturity, the selfed plants were individually harvested and seeds were catalogued and stored to ensure that the source of the seed was known.

Self-pollinated M₃ seed and Westar controls were analyzed in 10-seed bulk samples for fatty acid composition via gas chromatography. Statistical thresholds for each fatty acid component were established using a Z-distribution with a stringency level of 1 in 10,000. Mean and standard deviation values were determined from the non-mutagenized Westar control population in the field. The upper and lower statistical thresholds for each fatty acid were determined from the mean value of the population \pm the standard deviation, multiplied by the Z-distribution. Based on a population size of 10,000, the confidence interval is 99.99%.

The selected M₃ seeds were planted in the greenhouse along with Westar controls. The seed was sown in 4-inch pots containing Pro-Mix soil and the plants were maintained at 25°C/15°C, 14/10 hr day/night cycle in the greenhouse. At flowering, the terminal raceme was self-pollinated by bagging. At maturity, selfed M₄ seed was individually

harvested from each plant, labelled, and stored to ensure that the source of the seed was known.

The M_4 seed was analyzed in 10-seed bulk samples. Statistical thresholds for each fatty acid component were established from 259 control samples using a Z-distribution of 1 in 800. Selected M_4 lines were planted in a field trial in Carman, Manitoba in 3-meter rows with 6-inch spacing. Ten M_4 plants in each row were bagged for self-pollination. At maturity, the selfed plants were individually harvested and the open pollinated plants in the row were bulk harvested. The M_5 seed from single plant selections was analyzed in 10-seed bulk samples and the bulk row harvest in 50-seed bulk samples.

Selected M_5 lines were planted in the greenhouse along with Westar controls. The seed was grown as previously described. At flowering the terminal raceme was self-pollinated by bagging. At maturity, selfed M_6 seed was individually harvested from each plant and analyzed in 10-seed bulk samples for fatty acid composition.

Selected M_6 lines were entered into field trials in Eastern Idaho. The four trial locations were selected for the wide variability in growing conditions. The locations included Burley, Tetonia, Lamont and Shelley (Table 7). The lines were planted in four 3-meter rows with an 8-inch spacing, each plot was replicated four times. The planting design was determined using a Randomized Complete Block Designed. The commercial cultivar Westar was used as a check cultivar. At maturity the plots were harvested to determine yield. Yield of the entries in the trial was determined by taking the statistical average of the four replications. The Least Significant Difference Test was used to rank the entries in the randomized complete block design.

TABLE 7

Trial Locations for Selected Fatty Acid Mutants

<u>LOCATION</u>	<u>SITE CHARACTERIZATIONS</u>
BURLEY	Irrigated. Long season. High temperatures during flowering.
5 TETONIA	Dryland. Short season. Cool temperatures.
LAMONT	Dryland. Short season. Cool temperatures.
SHELLEY	Irrigated. Medium season. High temperatures during flowering.

To determine the fatty acid profile of entries, plants in each plot were bagged for self-pollination. The
 10 M₇ seed from single plants was analyzed for fatty acids in ten-seed bulk samples.

To determine the genetic relationships of the selected fatty acid mutants crosses were made. Flowers of M₆ or later generation mutations were used in crossing. F₁
 15 seed was harvested and analyzed for fatty acid composition to determine the mode of gene action. The F₁ progeny were planted in the greenhouse. The resulting plants were self-pollinated, the F₂ seed harvested and analyzed for fatty acid composition for allelism studies. The F₂ seed and
 20 parent line seed was planted in the greenhouse, individual plants were self-pollinated. The F₃ seed of individual plants was tested for fatty acid composition using 10-seed bulk samples as described previously.

In the analysis of some genetic relationships
 25 dihaploid populations were made from the microspores of the F₁ hybrids. Self-pollinated seed from dihaploid plants were analyzed for fatty acid analysis using methods described previously.

For chemical analysis, 10-seed bulk samples were
 30 hand ground with a glass rod in a 15-mL polypropylene tube

and extracted in 1.2 mL 0.25 N KOH in 1:1 ether/methanol. The sample was vortexed for 30 sec. and heated for 60 sec. in a 60°C water bath. Four mL of saturated NaCl and 2.4 mL of iso-octane were added, and the mixture was vortexed again. After phase separation, 600 µL of the upper organic phase were pipetted into individual vials and stored under nitrogen at -5°C. One µL samples were injected into a Supelco SP-2330 fused silica capillary column (0.25 mm ID, 30 M length, 0.20 µm df).

10 The gas chromatograph was set at 180°C for 5.5 minutes, then programmed for a 2°C/minute increase to 212°C, and held at this temperature for 1.5 minutes. Total run time was 23 minutes. Chromatography settings were: Column head pressure - 15 psi, Column flow (He) - 0.7 mL/min.,
15 Auxiliary and Column flow - 33 mL/min., Hydrogen flow - 33 mL/min., Air flow - 400 mL/min., Injector temperature - 250°C, Detector temperature - 300°C, Split vent - 1/15.

Table 8 describes the upper and lower statistical thresholds for each fatty acid of interest.

TABLE 8

Statistical Thresholds for Specific Fatty Acids
Derived from Control Westar Plantings

5	Genotype	Percent Fatty Acids				
		C _{16:0}	C _{18:0}	C _{18:1}	C _{18:2}	C _{18:3}
	M ₃ Generation (1 in 10,000 rejection rate)					
	Lower	3.3	1.4	--	13.2	5.3
	Upper	4.3	2.5	71.0	21.6	9.9
10	M ₄ Generation (1 in 800 rejection rate)					
	Lower	3.6	0.8	--	12.2	3.2
	Upper	6.3	3.1	76.0	32.4	9.9
	M ₅ Generation (1 in 755 rejection rate)					
	Lower	2.7	0.9	--	9.6	2.6
15	Upper	5.7	2.7	80.3	26.7	9.6

*Sats=Total Saturate Content

EXAMPLE 2

High Oleic Acid Canola Lines

In the studies of Example 1, at the M₃ generation,
20 31 lines exceeded the upper statistical threshold for oleic
acid ($\geq 71.0\%$). Line W7608.3 had 71.2% oleic acid. At the
M₄ generation, its selfed progeny (W7608.3.5, since
designated A129.5) continued to exceed the upper statistical
threshold for C_{18:1} with 78.8% oleic acid. M₅ seed of five
25 self-pollinated plants of line A129.5 (ATCC 40811) averaged
75.0% oleic acid. A single plant selection, A129.5.3 had
75.6% oleic acid. The fatty acid composition of this high
oleic acid mutant, which was stable under both field and
greenhouse conditions to the M₇ generation, is summarized in

Table 9. This line also stably maintained its mutant fatty acid composition to the M₇ generation in field trials in multiple locations. Over all locations the self-pollinated plants (A129) averaged 78.3% oleic acid. The fatty acid composition of the A129 for each Idaho trial location are summarized in Table 10. In multiple location replicated yield trials, A129 was not significantly different in yield from the parent cultivar Westar.

The canola oil of A129, after commercial processing, was found to have superior oxidative stability compared to Westar when measured by the Accelerated Oxygen Method (AOM), American Oil Chemists' Society Official Method Cd 12-57 for fat stability; Active Oxygen Method (revised 1989). The AOM of Westar was 18 AOM hours and for A129 was 30 AOM hours.

TABLE 9

Fatty Acid Composition of a High
Oleic Acid Canola Line Produced by Seed Mutagenesis

		Percent Fatty Acids				
Genotype	C _{16:0}	C _{18:0}	C _{18:1}	C _{18:2}	C _{18:3}	Sats
Westar	3.9	1.9	67.5	17.6	7.4	7.0
W7608.3 (M ₃)	3.9	2.4	71.2	12.7	6.1	7.6
W7608.3.5 (M ₄)	3.9	2.0	78.8	7.7	3.9	7.3
A129.5.3 (M ₅)	3.8	2.3	75.6	9.5	4.9	7.6

Sats=Total Saturate Content

TABLE 10

Fatty Acid Composition of a Mutant High Oleic Acid Line at Different Field Locations in Idaho

Percent Fatty Acids						
Location	C _{16:0}	C _{18:0}	C _{18:1}	C _{18:2}	C _{18:3}	Sats
Burley	3.3	2.1	77.5	8.1	6.0	6.5
Tetonia	3.5	3.4	77.8	6.5	4.7	8.5
Lamont	3.4	1.9	77.8	7.4	6.5	6.3
Shelley	3.3	2.6	80.0	5.7	4.5	7.7
Sats=Total Saturate Content						

The genetic relationship of the high oleic acid mutation A129 to other oleic desaturases was demonstrated in crosses made to commercial canola cultivars and a low linolenic acid mutation. A129 was crossed to the commercial cultivar Global (C_{16:0} - 4.5%, C_{18:0} - 1.5%, C_{18:1} - 62.9%, C_{18:2} - 20.0%, C_{18:3} - 7.3%). Approximately 200 F₂ individuals were analyzed for fatty acid composition. The results are summarized in Table 11. The segregation fit 1:2:1 ratio suggesting a single co-dominant gene controlled the inheritance of the high oleic acid phenotype.

TABLE 11

Genetic Studies of A129 X Global

Frequency			
Genotype	C _{18:0} Content (%)	Observed	Expected
od-od-	77.3	43	47
od-od+	71.7	106	94
od+od+	66.1	49	47

A cross between A129 and IMC 01, a low linolenic acid variety (C_{16:0} - 4.1%, C_{18:0} - 1.9%, C_{18:1} - 66.4%, C_{18:2} - 18.1%, C_{18:3} - 5.7%), was made to determine the inheritance of the oleic acid desaturase and linoleic acid desaturase.

In the F₁ hybrids both the oleic acid and linoleic acid desaturase genes approached the mid-parent values indicating a co-dominant gene actions. Fatty acid analysis of the F₂ individuals confirmed a 1:2:1:2:4:2:1:2:1 segregation of two independent, co-dominant genes (Table 12). A line was selected from the cross of A129 and IMC01 and designated as IMC130 (ATCC deposit no. 75446) as described in U.S. Patent Application No. 08/425,108, incorporated herein by reference.

TABLE 12

Genetic Studies of A129 X IMC 01

	Genotype	Ratio	Frequency	
			Observed	Expected
	od-od-ld-ld-	1	11	12
15	od-od-ld-ld+	2	30	24
	od-od-ld+ld+	1	10	12
	od-od+ld-ld-	2	25	24
	od-od+ld-ld+	4	54	47
	od-od+ld+ld+	2	18	24
20	od+od+ld-ld-	1	7	12
	od+od+ld-ld+	2	25	24
	od+od+ld+ld+	1	8	12

An additional high oleic acid line, designated A128.3, was also produced by the disclosed method. A 50-seed bulk analysis of this line showed the following fatty acid composition: C_{16:0} - 3.5%, C_{18:0} - 1.8%, C_{18:1} - 77.3%, C_{18:2} - 9.0%, C_{18:3} - 5.6%, FDA Sats - 5.3%, Total Sats - 6.4%.

This line also stably maintained its mutant fatty acid composition to the M₇ generation. In multiple locations replicated yield trials, A128 was not significantly different in yield from the parent cultivar Westar.

A129 was crossed to A128.3 for allelism studies. Fatty acid composition of the F₂ seed showed the two lines to be allelic. The mutational events in A129 and A128.3 although different in origin were in the same gene.

097704-01201
An additional high oleic acid line, designated M3028.-10 (ATCC 75026), was also produced by the disclosed method in Example 1. A 10-seed bulk analysis of this line showed the following fatty acid composition: C_{16:0} - 3.5%,
5 C_{18:0} - 1.8%, C_{18:1} - 77.3%, C_{18:2} - 9.0%, C_{18:3} - 5.6%, FDA Saturates - 5.3%, Total Saturates - 6.4%. In a single location replicated yield trial M3028.10 was not significantly different in yield from the parent cultivar Westar.

10 EXAMPLE 3

Low Linoleic Acid Canola

In the studies of Example 1, at the M₃ generation, 80 lines exceeded the lower statistical threshold for linoleic acid ($\leq 13.2\%$). Line W12638.8 had 9.4% linoleic
15 acid. At the M₄ and M₅ generations, its selfed progenies [W12638.8, since designated A133.1 (ATCC 40812)] continued to exceed the statistical threshold for low C_{18:2} with linoleic acid levels of 10.2% and 8.4%, respectively. The fatty acid composition of this low linoleic acid mutant,
20 which was stable to the M₅ generation under both field and greenhouse conditions, is summarized in Table 13. In multiple location replicated yield trials, A133 was not significantly different in yield from the parent cultivar Westar. An additional low linoleic acid line, designated
25 M3062.8 (ATCC 75025), was also produced by the disclosed method. A 10-seed bulk analysis of this line showed the following fatty acid composition: C_{16:0} - 3.8%, C_{18:0} - 2.3%, C_{18:1} - 77.1%, C_{18:2} - 8.9%, C_{18:3} - 4.3%, FDA Sats-6.1%. This line has also stably maintained its mutant fatty acid
30 composition in the field and greenhouse.

TABLE 13

Fatty Acid Composition of a Low
Linoleic Acid Canola Line Produced by Seed Mutagenesis

		Percent Fatty Acids				
	Genotype	C _{16:0}	C _{18:0}	C _{18:1}	C _{18:2}	Sats ^b
5	Westar	3.9	1.9	67.5	17.6	7.4
	W12638.8 (M ₃)	3.9	2.3	75.0	9.4	6.1
	W12638.8.1 (M ₄)	4.1	1.7	74.6	10.2	5.9
10	A133.1.8 (M ₅)	3.8	2.0	77.7	8.4	5.0
						7.0

^aLetter and numbers up to second decimal point indicate the plant line. Number after second decimal point indicates an individual plant.

^bSats=Total Saturate Content

EXAMPLE 4

Low Linolenic and Linoleic Acid Canola

In the studies of Example 1, at the M₃ generation, 57 lines exceeded the lower statistical threshold for linolenic acid ($\leq 5.3\%$). Line W14749.8 had 5.3% linolenic acid and 15.0% linoleic acid. At the M₄ and M₅ generations, its selfed progenies [W14749.8, since designated M3032 (ATCC 75021)] continued to exceed the statistical threshold for low C_{18:3} with linolenic acid levels of 2.7% and 2.3%, respectively, and for a low sum of linolenic and linoleic acids with totals of 11.8% and 12.5% respectively. The fatty acid composition of this low linolenic acid plus linoleic acid mutant, which was stable to the M₅ generation under both field and greenhouse conditions, is summarized in Table 14. In a single location replicated yield trial M3032

was not significantly different in yield from the parent cultivar (Westar).

TABLE 14

Fatty Acid Composition of a Low
Linolenic Acid Canola Line Produced by Seed Mutagenesis

Genotype	Percent Fatty Acids					Sats
	C _{16:0}	C _{18:0}	C _{18:1}	C _{18:2}	C _{18:3}	
Westar	3.9	1.9	67.5	17.6	7.4	7.0
W14749.8 (M ₃)	4.0	2.5	69.4	15.0	5.3	6.5
M3032.8 (M ₄)	3.9	2.4	77.9	9.1	2.7	6.4
M3032.1 (M ₅)	3.5	2.8	80.0	10.2	2.3	6.5
Sats=Total Saturate Content						

EXAMPLE 5

Canola Lines Q508 and Q4275

Seeds of the *B. napus* line IMC-129 were mutagenized with methyl N-nitrosoguanidine (MNNG). The MNNG treatment consisted of three parts: pre-soak, mutagen application, and wash. A 0.05M Sorenson's phosphate buffer was used to maintain pre-soak and mutagen treatment pH at 6.1. Two hundred seeds were treated at one time on filter paper (Whatman #3M) in a petri dish (100mm x 15mm). The seeds were pre-soaked in 15 mls of 0.05M Sorenson's buffer, pH 6.1, under continued agitation for two hours. At the end of the pre-soak period, the buffer was removed from the plate.

A 10mM concentration of MNNG in 0.05M Sorenson's buffer, pH 6.1, was prepared prior to use. Fifteen ml of 10m MNNG was added to the seeds in each plate. The seeds were incubated at 22°C±3°C in the dark under constant

agitation for four (4) hours. At the end of the incubation period, the mutagen solution was removed.

The seeds were washed with three changes of distilled water at 10 minute intervals. The fourth wash was for thirty minutes. This treatment regime produced an LD60 population.

Treated seeds were planted in standard greenhouse potting soil and placed into an environmentally controlled greenhouse. The plants were grown under sixteen hours of light. At flowering, the racemes were bagged to produce selfed seed. At maturity, the M2 seed was harvested. Each M2 line was given an identifying number. The entire MNNG-treated seed population was designated as the Q series.

Harvested M2 seeds was planted in the greenhouse. The growth conditions were maintained as previously described. The racemes were bagged at flowering for selfing. At maturity, the selfed M3 seed was harvested and analyzed for fatty acid composition. For each M3 seed line, approximately 10-15 seeds were analyzed in bulk as described in Example 1.

High oleic-low linoleic M3 lines were selected from the M3 population using a cutoff of >82% oleic acid and <5.0% linoleic. From the first 1600 M3 lines screened for fatty acid composition, Q508 was identified. The Q508 M3 generation was advanced to the M4 generation in the greenhouse. Table 15 shows the fatty acid composition of Q508 and IMC 129. The M4 selfed seed maintained the selected high oleic-low linoleic acid phenotype (Table 16).

TABLE 15

Fatty Acid Composition of A129 and High
Oleic Acid M3 Mutant Q508

Line #	16:0	18:0	18:1	18:2	18:3
5 A129 [*]	4.0	2.4	77.7	7.8	4.2
Q508	3.9	2.1	84.9	2.4	2.9

^{*}Fatty acid composition of A129 is the average of
50 self-pollinated plants grown with the M3 population

10 M₄ generation Q508 plants had poor agronomic
qualities in the field compared to Westar. Typical plants
were slow growing relative to Westar, lacked early
vegetative vigor, were short in stature, tended to be
chlorotic and had short pods. The yield of Q508 was very
low compared to Westar.

15 The M₄ generation Q508 plants in the greenhouse
tended to be reduced in vigor compared to Westar. However,
Q508 yields in the greenhouse were greater than Q508 yields
in the field.

TABLE 16

Fatty Acid Composition of Seed Oil
from Greenhouse-Grown Q508, IMC 129 and Westar.

Line	16:0	18:0	18:1	18:2	18:3	FDA Sats
20 IMC 129 ^a	4.0	2.4	77.7	7.8	4.2	6.4
25 Westar ^b	3.9	1.9	67.5	17.6	7.4	>5.8
Q508 ^c	3.9	2.1	84.9	2.4	2.9	6.0

^aAverage of 50 self-pollinated plants

^bData from Example 1

^cAverage of 50 self-pollinated plants

Nine other M4 high-oleic low-linoleic lines were also identified: Q3603, Q3733, Q4249, Q6284, Q6601, Q6761, Q7415, Q4275, and Q6676. Some of these lines had good agronomic characteristics and an elevated oleic acid level in seeds of about 80% to about 84%.

Q4275 was crossed to the variety Cyclone. After selfing for seven generations, mature seed was harvested from 93GS34-179, a progeny line of the Q4275 Cyclone cross. Referring to Table 17, fatty acid composition of a bulk seed sample shows that 93GS34 retained the seed fatty acid composition of Q4275. 93GS34-179 also maintained agronomically desirable characteristics.

After more than seven generations of selfing of Q4275, plants of Q4275, IMC 129 and 93GS34 were field grown during the summer season. The selections were tested in 4 replicated plots (5 feet X 20 feet) in a randomized block design. Plants were open pollinated. No selfed seed was produced. Each plot was harvested at maturity, and a sample of the bulk harvested seed from each line was analyzed for fatty acid composition as described above. The fatty acid compositions of the selected lines are shown in Table 17.

TABLE 17

Fatty Acid Composition of
Field Grown IMC 129, Q4275 and 93GS34 Seeds

Line	Fatty Acid Composition (%)					
	C _{16:0}	C _{18:0}	C _{18:1}	C _{18:2}	C _{18:3}	FDA Sats.
IMC 129	3.3	2.4	76.7	8.7	5.2	5.7
Q4275	3.7	3.1	82.1	4.0	3.5	6.8
93GS34-179	2.6	2.7	85.0	2.8	3.3	5.3

0971904-042904
The results shown in Table 17 show that Q4275 maintained the selected high oleic - low linoleic acid phenotype under field conditions. The agronomic characteristics of Q4275 plants were superior to those of Q508.

M₄ generation Q508 plants were crossed to a dihaploid selection of Westar, with Westar serving as the female parent. The resulting F₁ seed was termed the 92EF population. About 126 F₁ individuals that appeared to have better agronomic characteristics than the Q508 parent were selected for selfing. A portion of the F₂ seed from such individuals was replanted in the field. Each F₂ plant was selfed and a portion of the resulting F₃ seed was analyzed for fatty acid composition. The content of oleic acid in F₃ seed ranged from 59 to 79%. No high oleic (>80%) individuals were recovered with good agronomic type.

A portion of the F₂ seed of the 92EF population was planted in the greenhouse to analyze the genetics of the Q508 line. F₃ seed was analyzed from 380 F₂ individuals. The C_{18:1} levels of F₃ seed from the greenhouse experiment is depicted in Figure 1. The data were tested against the hypothesis that Q508 contains two mutant genes that are semi-dominant and additive: the original IMC 129 mutation as well as one additional mutation. The hypothesis also assumes that homozygous Q508 has greater than 85% oleic acid and homozygous Westar has 62-67% oleic acid. The possible genotypes at each gene in a cross of Q508 by Westar may be designated as:

AA = Westar Fad2^a

BB = Westar Fad2^b

aa = Q508 Fad2^{a-}

bb = Q508 Fad2^{b-}

Assuming independent segregation, a 1:4:6:4:1 ratio of phenotypes is expected. The phenotypes of heterozygous plants are assumed to be indistinguishable and, thus, the data were tested for fit to a 1:14:1 ratio of homozygous Westar: heterozygous plants: homozygous Q508.

Phenotypic	# of	
<u>Ratio</u>	<u>Westar Alleles</u>	<u>Genotype</u>
1	4	AABB (Westar)
4	3	AABb, AaBB, AABb, AaBB
10 6	2	AaBb, AAbb, AaBb, AaBb, aaBB, AaBb
4	1	Aabb, aaBb, Aabb, aaBb
1	0	aabb (Q508)

Using Chi-square analysis, the oleic acid data fit a 1:14:1 ratio. It was concluded that Q508 differs from Westar by two major genes that are semi-dominant and additive and that segregate independently. By comparison, the genotype of IMC 129 is aaBB.

The fatty acid composition of representative F3 individuals having greater than 85% oleic acid in seed oil is shown in Table 18. The levels of saturated fatty acids are seen to be decreased in such plants, compared to Westar.

TABLE 18

92EF F, Individuals with >85% C_{18:1} in Seed Oil

F3 Plant Identifier	Fatty Acid Composition (%)					
	C16:0	C18:0	C18:1	C18:2	C18:3	FDASA
+38068	3.401	1.582	85.452	2.134	3.615	4.983
+38156	3.388	1.379	85.434	2.143	3.701	4.767
+38171	3.588	1.511	85.289	2.367	3.425	5.099
+38181	3.75	1.16	85.312	2.968	3.819	4.977
+38182	3.529	0.985	85.905	2.614	3.926	4.56
+38191	3.364	1.039	85.737	2.869	4.039	4.459
+38196	3.557	1.182	85.054	2.962	4.252	4.739
+38202	3.554	1.105	86.091	2.651	3.721	4.713
+38220	3.093	1.16	86.421	1.931	3.514	4.314
+38236	3.308	1.349	85.425	2.37	3.605	4.718
+38408	3.617	1.607	85.34	2.33	3.562	5.224
+38427	3.494	1.454	85.924	2.206	3.289	4.948
+38533	3.64	1.319	85.962	2.715	3.516	4.959

EXAMPLE 6

Leaf and Root Fatty Acid Profiles of Canola
Lines IMC-129, Q508, and Westar

Plants of Q508, IMC 129 and Westar were grown in the greenhouse. Mature leaves, primary expanding leaves, petioles and roots were harvested at the 6-8 leaf stage,

09771904-012901
FOE2FO-406T2260

frozen in liquid nitrogen and stored at -70°C . Lipid extracts were analyzed by GLC as described in Example 1. The fatty acid profile data are shown in Table 19.

5 The data in Table 19 indicate that total leaf lipids in Q508 are higher in $\text{C}_{18:1}$ content than the $\text{C}_{18:2}$ plus $\text{C}_{18:3}$ content. The reverse is true for Westar and IMC 129. The difference in total leaf lipids between Q508 and IMC 129 is consistent with the hypothesis that a second Fad2 gene is mutated in Q508.

10 The $\text{C}_{16:3}$ content in the total lipid fraction was about the same for all three lines, suggesting that the plastid FadC gene product was not affected by the Q508 mutations. To confirm that the FadC gene was not mutated, chloroplast lipids were separated and analyzed. No changes
15 in chloroplast $\text{C}_{16:1}$, $\text{C}_{16:2}$ or $\text{C}_{16:3}$ fatty acids were detected in the three lines. The similarity in plastid leaf lipids among Q508, Westar and IMC 129 is consistent with the hypothesis that the second mutation in Q508 affects a microsomal Fad2 gene and not a plastid FadC gene.

TABLE 19

	MATURE LEAF			EXPANDING LEAF			PETIOLE			ROOT		
	West.	129	3Q508	West.	129	3Q508	West.	129	3Q508	West.	129	3Q508
16:0	12.1	11.9	10.1	16.4	16.1	11.3	21.7	23.5	11.9	21.1	21.9	12.0
16:1	0.8	0.6	1.1	0.7	0.6	1.1	1.0	1.3	1.4	-	-	-
16:2	2.3	2.2	2.0	2.8	3.1	2.8	1.8	2.2	1.8	-	-	-
16:3	14.7	15.0	14.3	6.3	5.4	6.9	5.7	4.6	5.7	-	-	-
18:0	2.2	1.6	1.2	2.5	2.8	1.5	3.7	4.0	1.6	3.6	2.9	2.5
18:1	2.8	4.9	16.7	3.8	8.3	38.0	4.9	12.9	46.9	3.5	6.1	68.8
18:2	12.6	11.5	6.8	13.3	13.8	4.9	20.7	18.3	5.2	28.0	30.4	4.4
18:3	50.6	50.3	46.0	54.2	50.0	33.5	40.4	33.2	25.3	43.8	38.7	12.3

EXAMPLE 7

Sequences of Mutant and Wild-Type Delta-12 Fatty Acid
Desaturases from *B. napus*

Primers specific for the FAD2 structural gene were used to clone the entire open reading frame (ORF) of the D and F delta-12 desaturase genes by reverse transcriptase polymerase chain reaction (RT-PCR). RNA from seeds of IMC 129, Q508 and Westar plants was isolated by standard methods and was used as template. The RT-amplified fragments were used for nucleotide sequence determination. The DNA sequence of each gene from each line was determined from both strands by standard dideoxy sequencing methods.

Sequence analysis revealed a G to A transversion at nucleotide 316 (from the translation initiation codon) of the D gene in both IMC 129 and Q508, compared to the sequence of Westar. The transversion changes the codon at this position from GAG to AAG and results in a non-conservative substitution of glutamic acid, an acidic residue, for lysine a basic residue. The presence of the

same mutation in both lines was expected since the Q508 line was derived from IMC 129. The same base change was also detected in Q508 and IMC 129 when RNA from leaf tissue was used as template.

5 The G to A mutation at nucleotide 316 was confirmed by sequencing several independent clones containing fragments amplified directly from genomic DNA of IMC 129 and Westar. These results eliminated the possibility of a rare mutation introduced during reverse transcription and PCR in
10 the RT-PCR protocol. It was concluded that the IMC 129 mutant is due to a single base transversion at nucleotide 316 in the coding region of the D gene of rapeseed microsomal delta 12-desaturase.

 A single base transition from T to A at nucleotide
15 515 of the F gene was detected in Q508 compared to the Westar sequence. The mutation changes the codon at this position from CTC to CAC, resulting in the non-conservative substitution of a ~~non-polar residue, leucine, for a polar residue, histidine~~^A, in the resulting gene product. No
20 mutations were found in the F gene sequence of IMC 129 compared to the F gene sequence of Westar.

 These data support the conclusion that a mutation in a delta-12 desaturase gene sequence results in alterations in the fatty acid profile of plants containing such a
25 mutated gene. Moreover, the data show that when a plant line or species contains two delta-12 desaturase loci, the fatty acid profile of an individual having two mutated loci differs from the fatty acid profile of an individual having one mutated locus.

30 The mutation in the D gene of IMC 129 and Q508 mapped to a region having a conserved amino acid motif (His-Xaa-Xaa-Xaa-His) found in cloned delta-12 and delta-15 membrane bound-desaturases (Table 20).

TABLE 20

Alignment of Amino Acid Sequences
of Cloned Canola Membrane Bound-Desaturases

Desaturase Gene	Sequence ^a	Position
Canola-fad2-D (mutant)	AHKCGH	109-114
Canola-Fad2-D	AHECGH	109-114
Canola-Fad2-F	AHECGH	109-114
Canola-FadC	<u>GHDCAH</u>	170-175
Canola-fad3 (mutant)	<u>GHKCGH</u>	94-99
Canola-Fad3	<u>GHDCGH</u>	94-99
Canola-FadD	<u>GHDCGH</u>	125-130

(FadD = Plastid delta 15, Fad3 = Microsomal delta-15),
(FadC = Plastid delta-12, Fad2 = Microsomal delta-12)

^a One letter amino acid code; conservative substitutions are underlined; non-conservative substitutions are in bold.

EXAMPLE 8

Transcription and Translation of Microsomal Delta-12
Fatty Acid Desaturases

Transcription in vivo was analyzed by RT-PCR analysis of stage II and stage III developing seeds and leaf tissue. The primers used to specifically amplify delta-12 desaturase F gene RNA from the indicated tissues were sense primer 5'-GGATATGATGATGGT^(Seq ID NO: 19)GAAAGA-3' and antisense primer 5'-TCTTTCACCATCATCATATCC-3'^(Seq ID NO: 20). The primers used to specifically amplify delta-12 desaturase D gene RNA from the indicated tissues were sense primer 5'-GTTATGAAGCAAAGAAGAAAC-3'^(Seq ID NO: 21) and antisense primer 5'-GTTTCTTCTTTGCTTCATAAC-3'^(Seq ID NO: 22). The results indicated that mRNA of both the D and F gene was expressed

Sub
C100
in seed and leaf tissues of IMC 129, Q508 and wild type Westar plants.

In vitro transcription and translation analysis showed that a peptide of about 46 kD was made. This is the expected size of both the D gene product and the F gene product, based on sum of the deduced amino acid sequence of each gene and the cotranslational addition of a microsomal membrane peptide.

These results rule out the possibility that non-sense or frameshift mutations, resulting in a truncated polypeptide gene product, are present in either the mutant D gene or the mutant F gene. The data, in conjunction with the data of Example 7, support the conclusion that the mutations in Q508 and IMC 129 are in delta-12 fatty acid desaturase structural genes encoding desaturase enzymes, rather than in regulatory genes.

EXAMPLE 9

Development of Gene-Specific PCR Markers

Based on the single base change in the mutant D gene of IMC 129 described in above, two 5' PCR primers were designed. The nucleotide sequence of the primers differed only in the base (G for Westar and A for IMC 129) at the 3' end. The primers allow one to distinguish between mutant fad2-D and wild-type Fad2-D alleles in a DNA-based PCR assay. Since there is only a single base difference in the 5' PCR primers, the PCR assay is very sensitive to the PCR conditions such as annealing temperature, cycle number, amount, and purity of DNA templates used. Assay conditions have been established that distinguish between the mutant gene and the wild type gene using genomic DNA from IMC 129 and wild type plants as templates. Conditions may be further optimized by varying PCR parameters, particularly

with variable crude DNA samples. A PCR assay distinguishing the single base mutation in IMC 129 from the wild type gene along with fatty acid composition analysis provides a means to simplify segregation and selection analysis of genetic crosses involving plants having a delta-12 fatty acid desaturase mutation.

EXAMPLE 10

Transformation with Mutant and Wild Type Fad3 Genes

B. napus cultivar Westar was transformed with mutant and wild type Fad3 genes to demonstrate that the mutant Fad3 gene for canola cytoplasmic linoleic desaturase delta-15 desaturase is nonfunctional. Transformation and regeneration were performed using disarmed *Agrobacterium tumefaciens* essentially following the procedure described in WO 94/11516.

Two disarmed *Agrobacterium* strains were engineered, each containing a Ti plasmid having the appropriate gene linked to a seed-specific promoter and a corresponding termination sequence. The first plasmid, pIMC110, was prepared by inserting into a disarmed Ti vector the full length wild type Fad3 gene in sense orientation (nucleotides 208 to 1336 of SEQ ID 6 in WO 93/11245), flanked by a napin promoter sequence positioned 5' to the Fad3 gene and a napin termination sequence positioned 3' to the Fad3 gene. The rapeseed napin promoter is described in EP 0255378.

The second plasmid, pIMC205, was prepared by inserting a mutated Fad3 gene in sense orientation into a disarmed Ti vector. The mutant sequence contained mutations at nucleotides 411 and 413 of the microsomal Fad3 gene described in WO93/11245, thus changing the sequence for codon 96 from GAC to AAG. The amino acid at codon 96 of the gene product was thereby changed from aspartic acid to lysine. See Table 20. A bean (*Phaseolus vulgaris*)

Sub
C8
could

phaseolin (7S seed storage protein) promoter fragment of 495 base pairs, starting with 5'-TGGTCTTTTGGT-3', was placed 5' to the mutant Fad3 gene and a phaseolin termination sequence was placed 3' to the mutant Fad3 gene. The phaseolin
5 sequence is described in Doyle et al., (1986) J. Biol. Chem. 261:9228-9238) and Slightom et al., (1983) Proc. Natl. Acad. Sci. USA 80:1897-1901.

The appropriate plasmids were engineered and transferred separately to Agrobacterium strain LEA4404.

10 Each engineered strain was used to infect 5 mm segments of hypocotyl explants from Westar seeds by cocultivation. Infected hypocotyls were transferred to callus medium and, subsequently, to regeneration medium. Once discernable
15 stems formed from the callus, shoots were excised and transferred to elongation medium. The elongated shoots were cut, dipped in Rootone™, rooted on an agar medium and transplanted to potting soil to obtain fertile T1 plants. T2 seeds were obtained by selfing the resulting T1 plants.

Fatty acid analysis of T2 seeds was carried out as
20 described above. The results are summarized in Table 21. Of the 40 transformants obtained using the pIMC110 plasmid, 17 plants demonstrated wild type fatty acid profiles and 16 demonstrated overexpression. A proportion of the transformants are expected to display an overexpression
25 phenotype when a functioning gene is transformed in sense orientation into plants.

Of the 307 transformed plants having the pIMC205 gene, none exhibited a fatty acid composition indicative of overexpression. This result indicates that the mutant fad3
30 gene product is non-functional, since some of the transformants would have exhibited an overexpression phenotype if the gene product were functional.

TABLE 21

Overexpression and Co-suppression Events in
Westar Populations Transformed with pIMC205 or pIMC110.

Construct	Number of Transforms	α -Linolenic Acid Range (%)	Overexpression Events (>10% linolenic)	Co-Suppression Events (<4.0% linolenic)	Wild Type Events
pIMC110	40	2.4 - 20.6	16	7	17
pIMC205	307	4.6 - 10.4	0	0	307

Fatty acid compositions of representative transformed plants are presented in Table 22. Lines 652-09 and 663-40 are representative of plants containing pIMC110 and exhibiting an overexpression and a co-suppression phenotype, respectively. Line 205-284 is representative of plants containing pIMC205 and having the mutant fad3 gene.

TABLE 22

Fatty Acid Composition of T2 Seed
From Westar Transformed With pIMC205 or pIMC110.

Line	Fatty Acid Composition (%)				
	C16:0	C18:0	C18:1	C18:2	C18:3
652-09 pIMC110 overexpression	4.7	3.3	65.6	8.1	14.8
663-40 pIMC110 co-suppression	4.9	2.1	62.5	23.2	3.6
205-284 pIMC205	3.7	1.8	68.8	15.9	6.7

EXAMPLE 11

Sequences of Wild Type and Mutant Fad2-D and Fad2-F

High molecular weight genomic DNA was isolated from leaves of Q4275 plants (Example 5). This DNA was used as template for amplification of Fad2-D and Fad2-F genes by polymerase chain reaction (PCR). PCR amplifications were carried out in a total volume of 100 μ l and contained 0.3 μ g genomic DNA, 200 μ M deoxyribonucleoside triphosphates, 3 mM

MgSO₄, 1-2 Units DNA polymerase and 1X Buffer (supplied by the DNA polymerase manufacturer). Cycle conditions were: 1 cycle for 1 min at 95°C, followed by 30 cycles of 1 min at 94°C, 2 min at 55°C and 3 min at 73°C.

5 The Fad2-D gene was amplified once using Elongase® (Gibco-BRL). PCR primers were: (SEQ ID NO: 23)
CAUCAUCAUCAUCTTCTTCGTAGGGTTCATCG and (SEQ ID NO: 24)
CUACUACUACUATCATAGAAGAGAAAGGTTTCAG_^ for the 5' and 3' ends of the gene, respectively.

10 The Fad2-F gene was independently amplified 4 times, twice with Elongase® and twice with Taq polymerase (Boehringer Mannheim). The PCR primers used were: (SEQ ID NO: 25)
5' CAUCAUCAUCAUATGGGTGCACGTGGAAGAA3' and (SEQ ID NO: 26)
5' CUACUACUACUATCTTTTACCATCATCATATCC3' for the 5' and 3' ends of the gene, respectively.

15 Amplified DNA products were resolved on an agarose gel, purified by JetSorb® and then annealed into pAMP1 (Gibco-BRL) via the (CAU)₄ and (CUA)₄ sequences at the ends of the primers, and transformed into *E. coli* DH5α.

20 The Fad2-D and Fad2-F inserts were sequenced on both strands with an ABI PRISM 310 automated sequencer (Perkin-Elmer) following the manufacturer's directions, using synthetic primers, AmpliTaq® DNA polymerase and dye terminator.

25 The Fad2-D gene was found to have an intron upstream of the ATG start codon. As expected, the coding sequence of the gene contained a G to A mutation at nucleotide 316, derived from IMC 129 (Fig. 2).

30 A single base transversion from G to A at nucleotide 908 was detected in the F gene sequence of the Q4275 amplified products, compared to the wild type F gene sequence (Fig. 2). This mutation changes the codon at amino acid 303 from GGA to GAA, resulting in the non-conservative

substitution of a glutamic acid residue for a glycine residue (Table 3 and Fig. 3). Expression of the mutant Q4275 Fad2-F delta-12 desaturase gene in plants alters the fatty acid composition, as described hereinabove.

5 To the extent not already indicated, it will be understood by those of ordinary skill in the art that any one of the various specific embodiments herein described and illustrated may be further modified to incorporate features shown in other of the specific embodiments.

10 The foregoing detailed description has been provided for a better understanding of the invention only and no unnecessary limitation should be understood therefrom as some modifications will be apparent to those skilled in the art without deviating from the spirit and scope of the
15 appended claims.

0971904-012901